

West Bengal State University
B.A./B.Sc./B.Com (Honours, Major, General) Examinations, 2015

PART - I

CHEMISTRY — HONOURS

Paper - II

Duration : 2 Hours]

[Full Marks : 50

The figures in the margin indicate full marks.

CEMAT-12-PA

Answer any two questions taking one from each Unit.

UNIT - I

1. a) The Maxwell's speed distribution is given as

$$P(c) = A c^2 e^{-mc^2/(2kT)}$$

What does $P(c)$ mean? Find the dimension of A .

Find $\left\langle \frac{1}{C} \right\rangle$ [Given : $\int_0^{\infty} x \exp(-\beta x^2) dx = \frac{1}{(2\beta)}$] 1 + 1 + 3

- b) What do you mean by 'collision diameter'? Derive a relation between mean free path (λ) and collision diameter of a gas. How does λ depend on T and P ? 1 + 3 + 1
- c) Apply the equipartition principle to calculate $\gamma (\approx C_p/C_v)$ for a linear, tri-atomic, ideal gas. What is the limiting value of γ when the number of atoms in the molecules becomes very large? 2 + 1
2. a) Obtain an expression for the distribution of translational kinetic energy from Maxwell's distribution of speed. How does it differ if the gas is changed from Helium to Argon? 4 + 1
- b) Calculate the number of binary collision per c.c. of $N_2(g)$ per second, at 2 atm, 30°C. The bond length of the gas molecules is 1.87 Å. 3

- c) Two flasks *A* and *B* have equal volumes. *A* is maintained at 300 K and *B* at 600 K. *A* contains hydrogen gas and *B* has equal mass of methane gas. Assuming ideal behaviour for both the gases, calculate (i) the ratio of mean free path in two containers [The collision diameter of methane may be assumed to be twice as that of hydrogen], (ii) the ratio of coefficient of viscosity in two flasks. 5

UNIT - II

3. a) Explain the term 'contact angle' and indicate explicitly the factors which govern its value when a liquid comes in contact with a solid in a gaseous atmosphere. 4
- b) At room temperature iodine is solid, bromine is liquid, chlorine and fluorine are gaseous. Can you explain this on the basis of intermolecular attraction? 2
- c) Write the unit and dimension of van der Waals constants '*a*' and '*b*'. 1 + 1
- d) A steel ball of density 7.9 g/c.c. having diameter of 4 mm is dropped into a column of liquid. It takes 55 seconds to fall through a distance of 1 m. Calculate the viscosity of the liquid. [Given, the density of the liquid is 1.10 g/c.c.] 3
- e) What do you mean by 'Boyle temperature'? 1
4. a) Give the graphical representation of a van der Waals gas explaining the different regions and parts. 3 + 2
Explain critical temperature (T_c) with reference to the curve.
- b) What is 'virial equation of state'? What does the second virial coefficient (B_{2v}) signify? 2
- c) What are viscosity and viscosity coefficient? Explain the unit of the coefficient. 3
- d) Derive Laplace's equation for excess pressure inside a spherical bubble, suspended in air. 2

CEMAT-12-PB

Answer any two questions taking one from each Unit.

UNIT - I

5. a) For a constant pressure process, $\Delta H = q_p$. Does it follow that q_p is a state function? 2
- b) Why does C_p exceeds C_v for an ideal gas? Give a molecular explanation. 2
- c) Show that $C_p - C_v = \frac{T\alpha^2 V}{\kappa}$, where α = coefficient of volume expansion; κ = coefficient of compressibility. 4
- d) At 25°C the coefficient of thermal expansion of water is $\alpha = 2.07 \times 10^{-4} \text{ K}^{-1}$ and the density is 0.9970 g/cm³. If the temperature of 200 g of water is raised from 25°C to 50°C under a constant pressure of 101 kPa, calculate w . Calculate also q , ΔH and ΔU . (Given, $C_p = 75.30 \text{ JK}^{-1} \text{ mol}^{-1}$) 3
- e) A spontaneous polymerisation must be an exothermic process." Justify or criticize. 2
6. a) " C_v is independent of pressure for a gas obeying equation of state $P(V - nb) = nRT$." Justify or criticize. 2
- b) Show the equivalence of Kelvin-Planck and Clausius statements of the second law of thermodynamics. 4
- c) Show that for a van der Waals gas, $\frac{\alpha}{\kappa} = \frac{R}{V - b}$ where, α = coefficient of volume expansion; κ = coefficient of compressibility. 3
- d) Draw a Carnot cycle in H vs T and U vs S diagram. 2
- e) A Carnot cycle heat engine does 2.50 kJ of work per cycle and has an efficiency of 45%. Find the temperatures of the two reservoirs between which the Carnot engine is operating. 2

UNIT - II

7. a) For a consecutive reaction $A \rightarrow B \rightarrow C$, obtain a relation for the time at which the concentration of B will be maximum in terms of rate constants of individual steps. Assume all the reaction steps are first order. 4
- b) "Second order reaction is never completed." Justify or criticize. 2

- c) The rate constant of a reaction is experimentally found to decrease with increase in temperature. Is there any contradiction with Arrhenius rate equation? 2
- d) Show that if A undergoes two simultaneous reactions (parallel reactions) to produce B and C according to the reactions $A \rightarrow B$ and $A \rightarrow C$, then the observed activation energy (E_a) for the disappearance of A is given by the equation $E_a = \frac{k_1 E_1 + k_2 E_2}{k_1 + k_2}$, where k_1 and k_2 are the first order rate constants of the individual step and E_1 and E_2 are activation energies of the two steps. 4
8. a) "Unimolecular reactions are not always first order." Justify it using Lindemann's theory. 3
- b) "Zero order reaction must be a multi-step reaction." Justify or criticize. 2
- c) Consider a reaction $A \rightarrow$ Products, which is one-half order with respect to A .
- Derive the integrated form of rate equation.
 - Write the unit of rate constant (k).
 - Decide what function should be plotted to determine the rate constant. Also show the plot. $2 + \frac{1}{2} + (1 + \frac{1}{2})$
- d) "Activation energy of a reaction can never be zero or negative." Justify or criticize. 3